

W(5): Wobbling mode in the framework of the X(5) model

Dennis Bonatsos^{#1}, D. Lenis^{#2}, D. Petrellis^{#3}, P. A. Terziev^{†4}

[#] Institute of Nuclear Physics, N.C.S.R. “Demokritos”
GR-15310 Aghia Paraskevi, Attiki, Greece

[†] Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences
72 Tzarigrad Road, BG-1784 Sofia, Bulgaria

Abstract

Using in the Bohr Hamiltonian the approximations leading to the Bohr and Mottelson description of wobbling motion in even nuclei, a W(5) model for wobbling bands, coexisting with a X(5) ground state band, is obtained. Separation of variables is achieved by assuming that the relevant potential has a sharp minimum at γ_0 , which is the only parameter entering in the spectra and B(E2) transition rates (up to overall scale factors). B(E2) transition rates exhibit the features expected in the wobbling case, while the spectrum for $\gamma = 20^\circ$ is in good agreement with experimental data for ^{156}Dy .

1. Introduction

Nuclear wobbling motion [1] is expected to occur for triaxial nuclei at high angular momenta, when the angular momentum is aligned with the axis corresponding to the largest moment of inertia, a situation which classically corresponds to simple rotation without precession of the axes. Although wobbling motion was initially introduced for even nuclei [1], it has been seen experimentally up to now (and only recently) only in odd nuclei (^{163}Lu [2, 3, 4], ^{165}Lu [5], ^{167}Lu [6]). Detailed theoretical works have been performed in the cranked shell model plus random phase approximation [7, 8, 9], as well as in the particle-rotor model [10, 11], which naturally contain free parameters.

In the present work we attempt a nearly parameter-free (up to overall scale factors) description of wobbling in even nuclei, following the methods developed in the E(5) [12], X(5) [13], Y(5) [14], and Z(5) [15] models, which correspond to the U(5)–O(6), U(5)–SU(3), axial–triaxial, and prolate–oblate shape phase transitions respectively. Furthermore, the wobbling nucleus is assumed to possess a relatively rigid triaxial shape, as in Refs. [16, 17, 18], with the potential having a sharp minimum at $\gamma = \gamma_0$. γ_0 is the only free parameter entering in the problem. It will be seen, however, that the results are changing very little with γ_0 within the region of interest. The path we follow is described here:

1) We assume that the ground state band (gsb), which should be Yrast at low angular momentum L , is axial, characterized by $\gamma_0 = 0$. We then use for this purpose the X(5) gsb, which is indeed derived from the original Bohr Hamiltonian [19] after approximately separating variables for $\gamma = 0$ [13].

¹e-mail: bonat@inp.demokritos.gr

²e-mail: lenis@inp.demokritos.gr

³e-mail: petrellis@inp.demokritos.gr

⁴e-mail: terziev@inrne.bas.bg

2) We assume (as in Ref. [20]) that triaxiality should appear at higher L . Starting then from the original Bohr Hamiltonian, we approximately separate variables following the steps of Bohr and Mottelson [1] in the definitive description of wobbling and keeping γ close to γ_0 . The resulting model, in which only γ_0 appears as a parameter, we call W(5). The spectrum of W(5) is measured from the ground state of X(5) and normalized to the first excited state of the gsb of X(5), in order to be directly comparable to the X(5) spectrum.

3) The $n_w = 0$ band of W(5) (where n_w is the number of wobbling phonons [1]) is found to cross the gsb of X(5) at certain L , depending (very weakly within the region of interest) on γ_0 . Thus the $n_w = 0$ band of W(5) becomes Yrast beyond some specific L . Bands with $n_w = 1, 2, \dots$ exist at higher energies.

4) The $n_w = 0, 1, 2$ bands of W(5) are connected by intraband and interband B(E2) transitions which exhibit the characteristic features expected in the case of wobbling [21].

It is clear that the W(5) model should be tested against experiment in nuclei of which the gsb at low L appears to be close to X(5). A summary of such nuclei in the rare earth region is given in Ref. [22]. It is indeed seen that existing experimental spectra on ^{156}Dy [23] correspond very well to the $n_w = 0$ and $n_w = 1$ bands of the W(5) model for $\gamma_0 = 20^\circ$.

In Sections 2 and 3 of the present work the β -part and the γ -part of the W(5) spectrum are derived respectively, while B(E2) transition rates are studied in Section 4. Numerical results are presented in Section 5, while Section 6 contains a brief comparison to experiment and in Section 7 discussion of the present results and plans for further work are given.

2. The β -part of the spectrum

The original Bohr Hamiltonian [19] is

$$H = -\frac{\hbar^2}{2B} \left[\frac{1}{\beta^4} \frac{\partial}{\partial \beta} \beta^4 \frac{\partial}{\partial \beta} + \frac{1}{\beta^2 \sin 3\gamma} \frac{\partial}{\partial \gamma} \sin 3\gamma \frac{\partial}{\partial \gamma} - \frac{1}{4\beta^2} \sum_{k=1,2,3} \frac{Q_k^2}{\sin^2 \left(\gamma - \frac{2}{3}\pi k \right)} \right] + V(\beta, \gamma), \quad (1)$$

where β and γ are the usual collective coordinates, while Q_k ($k = 1, 2, 3$) are the components of angular momentum and B is the mass parameter.

Introducing [13] reduced energies $\epsilon = 2BE/\hbar^2$ and reduced potentials $u = 2BV/\hbar^2$, one aims at an approximate separation of variables by assuming that the reduced potential can be separated into two terms, one depending on β and the other depending on γ , i.e. $u(\beta, \gamma) = u(\beta) + u(\gamma)$.

In the X(5) model [13], approximate separation of variables is achieved by assuming that the potential $u(\gamma)$ has a minimum around $\gamma_0 = 0$, guaranteeing that K , the projection of angular momentum on the body-fixed $\hat{z}'$ -axis, is a good quantum number. One then seeks solutions of the relevant Schrödinger equation having the form $\Psi(\beta, \gamma, \theta_i) = \phi_K^L(\beta, \gamma) \mathcal{D}_{M,K}^L(\theta_i)$, where θ_i ($i = 1, 2, 3$) are the Euler angles, $\mathcal{D}(\theta_i)$ denote Wigner functions of them, while L and M are the eigenvalues of angular momentum and the eigenvalues of the projection of angular momentum on the laboratory-fixed $\hat{z}$ -axis respectively. In the case in which $u(\beta)$ is an infinite well potential

$$u(\beta) = \begin{cases} 0 & \text{if } \beta \leq \beta_W \\ \infty & \text{for } \beta > \beta_W \end{cases}, \quad (2)$$

the relevant differential equation is solved exactly, the corresponding eigenvalues being

$$\epsilon_{\beta;s,\nu} = (k_{s,\nu})^2, \quad k_{s,\nu} = \frac{x_{s,\nu}}{\beta_W}, \quad (3)$$

where $x_{s,\nu}$ is the s -th zero of the Bessel function $J_\nu(k_{s,\nu}\beta)$, with [13, 24]

$$\nu = \left(\frac{L(L+1) - K^2}{3} + \frac{9}{4} \right)^{1/2}. \quad (4)$$

while the relevant eigenfunctions are

$$\xi_{s,\nu}(\beta) = c_{s,\nu} \beta^{-3/2} J_\nu(k_{s,\nu}\beta), \quad (5)$$

where $c_{s,\nu}$ are normalization constants.

In the Z(5) model [15], approximate separation of variables is achieved by assuming that the potential $u(\gamma)$ has a minimum around $\gamma_0 = \pi/6$, guaranteeing [25] that α , the projection of angular momentum on the body-fixed $\hat{x}'$ -axis, is a good quantum number. One then seeks solutions of the relevant Schrödinger equation having the form $\Psi(\beta, \gamma, \theta_i) = \xi_{L,\alpha}(\beta) \eta(\gamma) \mathcal{D}_{M,\alpha}^L(\theta_i)$. In the case of $u(\beta)$ being an infinite well potential (Eq. (2)) the relevant differential equation is solved exactly, the eigenvalues and eigenfunctions having the form given in Eqs. (3) and (5) respectively, with [15]

$$\nu = \frac{\sqrt{4L(L+1) - 3\alpha^2 + 9}}{2} = \frac{\sqrt{L(L+4) + 3n_w(2L - n_w) + 9}}{2}, \quad (6)$$

where $n_w = L - \alpha$ is the wobbling quantum number [25].

The X(5) and Z(5) solutions, briefly reviewed above, are obtained for specific values of γ_0 (0, $\pi/6$ respectively), and are valid for any value of the angular momentum L . A different approximate solution, which for brevity we are going to call W(5), can be obtained by following the steps of Bohr and Mottelson [1] for the description of wobbling motion. This solution will be obtained for a range of γ_0 values, but it will be valid only for large values of the angular momentum L , which is supposed to be aligned along the axis corresponding to the largest moment of inertia.

Using in Eq. (1) the definitions

$$A_k = \frac{1}{\sin^2\left(\gamma - \frac{2\pi}{3}k\right)}, \quad k = 1, 2, 3 \quad (7)$$

one sees that in the region $0 < \gamma < \pi/6$ one has $A_1 < A_2 < A_3$. Therefore the largest moment of inertia corresponds to $k = 1$. In what follows we are going to restrict ourselves to the $0 < \gamma < \pi/6$ region.

For large angular momenta L aligned along the $k = 1$ axis, following Bohr and Mottelson [1] one can see that the eigenvalues of the $A_1 Q_1^2 + A_2 Q_2^2 + A_3 Q_3^2$ term in Eq. (1) take the form

$$\varepsilon(n_w, L) = A_1 L(L+1) + 2A_1 L \left(n_w + \frac{1}{2} \right) A_w, \quad (8)$$

with

$$A_w = \sqrt{\left(\frac{A_2}{A_1} - 1 \right) \left(\frac{A_3}{A_1} - 1 \right)}, \quad (9)$$

where n_w is the number of the wobbling excitation quanta, for which the approximate (in the present case) relation $n_w = L - \alpha$ (where α is the projection of angular momentum on the $k = 1$ body-fixed axis, as before) holds. Since $\alpha \approx L$ and L is a good quantum number, α can be approximately treated as a good quantum number, too.

It should be remembered that the approximations carried out in Ref. [1] are valid for

$$L \gg \left(n_w + \frac{1}{2}\right) \frac{1}{A_w} \left(\frac{A_2 + A_3}{A_1} - 2\right) \equiv L_0, \quad (10)$$

where A_w is given by Eq. (9).

Using this result in the Schrödinger equation corresponding to the Hamiltonian of Eq. (1), one can separate it into two equations

$$\left[-\frac{1}{\beta^4} \frac{\partial}{\partial \beta} \beta^4 \frac{\partial}{\partial \beta} + \frac{1}{4\beta^2} A_1 \left(L(L+1) + 2L \left(n_w + \frac{1}{2}\right) A_w\right) + u(\beta)\right] \xi_{L,n_w}(\beta) = \epsilon_\beta \xi_{L,n_w}(\beta), \quad (11)$$

$$\left[-\frac{1}{\langle \beta^2 \rangle \sin 3\gamma} \frac{\partial}{\partial \gamma} \sin 3\gamma \frac{\partial}{\partial \gamma} + u(\gamma)\right] \eta(\gamma) = \epsilon_\gamma \eta(\gamma), \quad (12)$$

where $\langle \beta^2 \rangle$ is the average of β^2 over $\xi(\beta)$, and $\epsilon = \epsilon_\beta + \epsilon_\gamma$. Here we assume, as in Refs. [16, 17, 18], that the potential $u(\gamma)$ has a deep minimum at $\gamma = \gamma_0$, and that the variable γ remains “frozen” at the value γ_0 in A_1 and A_w , appearing in Eq. (11). Furthermore, the $\langle \beta^2 \rangle$ term introduces a “hidden” dependence on s , L , and n_w in Eq. (12). The approximate separation of the β and γ variables is achieved by considering an adiabatic approximation, as in the X(5) case [13, 24].

The total wave function should have the form

$$\Psi(\beta, \gamma, \theta_i) = \xi_{L,n_w}(\beta) \eta(\gamma) \mathcal{D}_{M,\alpha}^L(\theta_i), \quad (13)$$

with $\alpha = L - n_w$.

In the case in which $u(\beta)$ is the infinite well potential of Eq. (2), one can use the transformation [13] $\tilde{\xi}(\beta) = \beta^{3/2} \xi(\beta)$, as well as the definitions [13] $\epsilon_\beta = k_\beta^2$, $z = \beta k_\beta$, in order to bring Eq. (11) into the form of a Bessel equation

$$\frac{d^2 \tilde{\xi}}{dz^2} + \frac{1}{z} \frac{d\tilde{\xi}}{dz} + \left[1 - \frac{\nu^2}{z^2}\right] \tilde{\xi} = 0, \quad (14)$$

with

$$\nu = \frac{\sqrt{A_1 L(L+1) + 2A_1 L \left(n_w + \frac{1}{2}\right) A_w + 9}}{2}. \quad (15)$$

Then the boundary condition $\tilde{\xi}(\beta_W) = 0$ determines the spectrum

$$\epsilon_{\beta;s,\nu} = \epsilon_{\beta;s,n_w,L} = (k_{s,\nu})^2, \quad k_{s,\nu} = \frac{x_{s,\nu}}{\beta_W}, \quad (16)$$

and the eigenfunctions

$$\xi_{s,\nu}(\beta) = \xi_{s,n_w,L}(\beta) = \xi_{s,\alpha,L}(\beta) = c_{s,\nu} \beta^{-3/2} J_\nu(k_{s,\nu} \beta), \quad (17)$$

where $x_{s,\nu}$ is the s th zero of the Bessel function $J_\nu(z)$, while the normalization constants $c_{s,\nu}$ are determined from the normalization condition $\int_0^\infty \beta^4 \xi_{s,\nu}^2(\beta) d\beta = 1$. The notation for the roots has been kept the same as in Ref. [13], while for the energies the notation $E_{s,n_w,L}$ will be used. The lowest band corresponds to $s = 1$, $n_w = 0$ with $L = 0, 2, 4, \dots$, while the next bands are $s = 1$, $n_w = 1$ with $L = 1, 3, 5, \dots$, and $s = 1$, $n_w = 2$ with $L = 2, 4, 6, \dots$ [25].

In the special case of $\gamma_0 = \pi/6$ one can easily see from Eqs. (7), (9) that $A_1 = 1$, $A_2 = A_3 = 4$, $A_w = 3$. Then Eq. (15) takes the form $\nu = \frac{1}{2}\sqrt{L(L+4) + 6n_wL + 9}$, which is in agreement with the corresponding Z(5) expression of Eq. (6) up to terms of order n_w^2 .

3. The γ -part of the spectrum

The γ -part of the spectrum is obtained from Eq. (12), which can be simply rewritten as

$$\left[-\frac{1}{\langle \beta^2 \rangle} \left(\frac{\partial^2}{\partial \gamma^2} + 3 \frac{\cos 3\gamma}{\sin 3\gamma} \frac{\partial}{\partial \gamma} \right) + u(\gamma) \right] \eta(\gamma) = \epsilon_\gamma \eta(\gamma). \quad (18)$$

We consider a harmonic oscillator potential having a sharp minimum at $\gamma = \gamma_0$ ($0 < \gamma_0 \leq \pi/6$), i.e.

$$u(\gamma) = \frac{1}{2}c(\gamma - \gamma_0)^2 = \frac{1}{2}c\tilde{\gamma}^2, \quad \tilde{\gamma} = \gamma - \gamma_0. \quad (19)$$

The minimum is sharp as long as the constant c is taken to be sufficiently large.

Since we consider only small oscillations around γ_0 , the above equation can be brought into the form

$$\left[-\frac{\partial^2}{\partial \tilde{\gamma}^2} - 3 \cot 3\gamma_0 \frac{\partial}{\partial \tilde{\gamma}} + \frac{1}{2}c\langle \beta^2 \rangle \tilde{\gamma}^2 \right] \eta(\tilde{\gamma}) = \epsilon_{\tilde{\gamma}} \langle \beta^2 \rangle \eta(\tilde{\gamma}). \quad (20)$$

Using

$$\eta(\tilde{\gamma}) = \tilde{\eta}(\tilde{\gamma}) e^{-3(\cot 3\gamma_0)\tilde{\gamma}/2}, \quad (21)$$

Eq. (20) is brought into the form

$$\left(-\frac{\partial^2}{\partial \tilde{\gamma}^2} + \frac{1}{2}c\langle \beta^2 \rangle \tilde{\gamma}^2 \right) \tilde{\eta}(\tilde{\gamma}) = \left(\epsilon_{\tilde{\gamma}} \langle \beta^2 \rangle - \frac{9}{4}(\cot 3\gamma_0)^2 \right) \tilde{\eta}(\tilde{\gamma}), \quad (22)$$

which is a simple harmonic oscillator equation with energy eigenvalues

$$\tilde{\epsilon}_{\tilde{\gamma}} = \epsilon_{\tilde{\gamma}} \langle \beta^2 \rangle - \frac{9}{4}(\cot 3\gamma_0)^2 = \sqrt{2c\langle \beta^2 \rangle} \left(n_{\tilde{\gamma}} + \frac{1}{2} \right), \quad n_{\tilde{\gamma}} = 0, 1, 2, \dots \quad (23)$$

and eigenfunctions

$$\tilde{\eta}_{n_{\tilde{\gamma}}}(\tilde{\gamma}) = N_{n_{\tilde{\gamma}}} H_{n_{\tilde{\gamma}}}(b\tilde{\gamma}) e^{-b^2\tilde{\gamma}^2/2}, \quad b = \left(\frac{c\langle \beta^2 \rangle}{2} \right)^{1/4}, \quad (24)$$

with normalization constant

$$N_{n_{\tilde{\gamma}}} = \sqrt{\frac{b}{\sqrt{\pi} 2^{n_{\tilde{\gamma}}} n_{\tilde{\gamma}}!}}. \quad (25)$$

Similar potentials and solutions in the γ -variable have been considered in [18, 19].

The total energy in the case of the W(5) model is then

$$E(s, n_w, L, n_{\tilde{\gamma}}) = E_0 + A(x_{s,\nu})^2 + Bn_{\tilde{\gamma}}. \quad (26)$$

It should be noticed at this point that the γ -part of the spectrum is treated here only to lowest order approximation. Although this suffices for the purpose of the present work, in which only states with $n_{\tilde{\gamma}} = 0$ are considered, higher approximations might be necessary for the detailed study of states with $n_{\tilde{\gamma}} \neq 0$.

4. B(E2) transition rates

As in the Z(5) case, the quadrupole operator is given by

$$T_{\mu}^{(E2)} = t\beta \left[\mathcal{D}_{\mu,0}^{(2)}(\theta_i) \cos\left(\gamma - \frac{2\pi}{3}\right) + \frac{1}{\sqrt{2}}(\mathcal{D}_{\mu,2}^{(2)}(\theta_i) + \mathcal{D}_{\mu,-2}^{(2)}(\theta_i)) \sin\left(\gamma - \frac{2\pi}{3}\right) \right], \quad (27)$$

where t is a scale factor, while in the Wigner functions the quantum number α appears next to μ , and the quantity $\gamma - 2\pi/3$ in the trigonometric functions is obtained from $\gamma - 2\pi k/3$ for $k = 1$, since in the present case the projection α along the body-fixed $\hat{x}'$ -axis is used. Eq. (27) is equivalent to Eq. (3.6) of Ref. [7], up to a sign convention for γ . This indicates that the results of the present work correspond to region 2 ($-60^\circ < \gamma < 0^\circ$) [7] in the Lund convention. However, in the present case only the region $0^\circ < \gamma < 30^\circ$ is covered, because of the assumptions made in Section 2.

B(E2) transition rates are given by

$$B(E2; L_i \alpha_i \rightarrow L_f \alpha_f) = \frac{5}{16\pi} \frac{|\langle L_f \alpha_f || T^{(E2)} || L_i \alpha_i \rangle|^2}{(2L_i + 1)}. \quad (28)$$

The symmetrized wave function reads

$$\Psi(\beta, \gamma, \theta_i) = \xi_{s,n_w,L}(\beta) \eta(\tilde{\gamma}) \sqrt{\frac{2L+1}{16\pi^2(1+\delta_{\alpha,0})}} (\mathcal{D}_{\mu,\alpha}^{(L)} + (-1)^L \mathcal{D}_{\mu,-\alpha}^{(L)}), \quad (29)$$

where the normalization factor occurs from the standard integrals involving two Wigner functions [26] and is the same as in Ref. [25]. α has to be an even integer [25], while for $\alpha = 0$ it is clear that only even values of L are allowed, since the symmetrized wave function is vanishing otherwise.

In the calculation of the matrix elements of Eq. (28) the integral over γ leads to unity [because of the normalization of $\eta(\tilde{\gamma})$], the integral over β takes the form

$$I_{\beta}(s_i, L_i, \alpha_i, s_f, L_f, \alpha_f) = \int \beta \xi_{s_i, \alpha_i, L_i}(\beta) \xi_{s_f, \alpha_f, L_f}(\beta) \beta^4 d\beta, \quad (30)$$

where the β factor comes from Eq. (27), and the β^4 factor comes from the volume element [19], while the integral over the angles is calculated using the standard integrals involving three Wigner functions [26]. The separation of the integrals occurs because $\eta(\tilde{\gamma})$ does not depend on α or n_w , while in $\xi(\beta)$ only even values of α appear. The final result reads

$$B(E2; L_i \alpha_i \rightarrow L_f \alpha_f) = \frac{5}{16\pi} t^2 \frac{1}{(1+\delta_{\alpha_i,0})(1+\delta_{\alpha_f,0})}$$

$$\begin{aligned}
& \left[\cos \left(\gamma - \frac{2\pi}{3} \right) \left\{ (L_i 2L_f | \alpha_i 0 \alpha_f) + (-1)^{L_i} (L_i 2L_f | -\alpha_i 0 \alpha_f) \right\} \right. \\
& \left. + \frac{1}{\sqrt{2}} \sin \left(\gamma - \frac{2\pi}{3} \right) \left\{ (L_i 2L_f | \alpha_i 2 \alpha_f) + (L_i 2L_f | \alpha_i - 2 \alpha_f) + (-1)^{L_i} (L_i 2L_f | -\alpha_i 2 \alpha_f) \right\} \right]^2 \\
& I_\beta^2(s_i, L_i, \alpha_i, s_f, L_f, \alpha_f).
\end{aligned} \tag{31}$$

One can easily see that the Clebsch–Gordan coefficients (CGCs) appearing in this equation impose a $\Delta\alpha = 0, \pm 2$ selection rule. Indeed, the first CGC is nonvanishing only for $\alpha_i = \alpha_f$, while the third CGC is nonvanishing only if $\alpha_i + 2 = \alpha_f$, and the fourth CGC is nonvanishing only if $\alpha_i - 2 = \alpha_f$. The second and fifth CGCs are nonvanishing only if $\alpha_i + \alpha_f = 0$ and $\alpha_i + \alpha_f = 2$ respectively, which can be valid only in a few special cases.

It is worth remarking that the first CGC in Eq. (31) allows for nonvanishing quadrupole moments, in contrast to the $Z(5)$ case [15], where quadrupole moments vanish up to this order.

5. Numerical results

For low angular momentum the nucleus is expected to have $\gamma_0 = 0$. As angular momentum rises, at some point the nucleus will “jump” (as in Ref. [20]) to the large L limit corresponding to wobbling motion. As a consequence, the ground state band (gsb) of the nucleus should correspond to the gsb of $X(5)$. The $X(5)$ gsb should be the Yrast band up to some value of L , beyond which the $n_w = 0$ wobbling band should become Yrast, while additional wobbling bands with $n_w = 1, 2, \dots$ should be seen further up in energy.

It is therefore reasonable to measure all energies from the ground state of $X(5)$ and normalize them to the lowest excitation energy of $X(5)$. Therefore for the wobbling levels the ratios

$$E_{s,n_w,L}^{W(5)} = \frac{\epsilon_{\beta,s,n_w,L}^{W(5)} - \epsilon_{\beta,s=1,L=0}^{X(5)}}{\epsilon_{\beta,s=1,L=2}^{X(5)} - \epsilon_{\beta,s=1,L=0}^{X(5)}} \tag{32}$$

will be used, while for the $X(5)$ levels the ratios

$$E_{s,L}^{X(5)} = \frac{\epsilon_{\beta,s,L}^{X(5)} - \epsilon_{\beta,s=1,L=0}^{X(5)}}{\epsilon_{\beta,s=1,L=2}^{X(5)} - \epsilon_{\beta,s=1,L=0}^{X(5)}} \tag{33}$$

will be used.

As we have seen in Section 2, the results depend on γ_0 , the value of γ at which the relevant potential is supposed to have a sharp minimum and at which the variable γ is “frozen” in the β -equation (Eq. (11)). In what follows we are going to focus attention on $\gamma_0 = 15^\circ$, which lies midway between the axial ($\gamma_0 = 0$) and maximally triaxial ($\gamma_0 = 30^\circ$) cases, and on $\gamma_0 = 20^\circ$, which has been found of interest [4] in the framework of “Ultimate Cranker” [27] calculations.

Spectra for the lowest wobbling bands for $\gamma = 15^\circ$ and $\gamma = 20^\circ$ are shown in Table 1, where the gsb of $X(5)$ is also shown for comparison. In Fig. 1 several levels of the $n_w = 0$ and $n_w = 1$ bands are plotted as a function of γ_0 . It is seen that the γ_0 -dependence of the energy levels is rather flat within the region of interest ($10^\circ < \gamma_0 < 30^\circ$), therefore the two γ_0 values shown in Table 1 suffice.

In Table 1 the restrictions imposed by the condition of Eq. (10) are reminded by reporting the appropriate L_0 values and by putting in parentheses the energies of the levels below this limit.

In Table 1 we remark that the $n_w = 0$ band with $\gamma_0 = 15^\circ$ crosses the X(5) gsb above $L = 12$ and becomes Yrast from $L = 14$ up, while the $n_w = 0$ band with $\gamma_0 = 20^\circ$ crosses the X(5) gsb above $L = 8$ and becomes Yrast from $L = 10$ up. The angular momentum L at which the bandcrossing of the X(5) gsb and the $n_w = 0$ band occurs does not depend on any free parameter, but only on γ_0 , the relevant dependence being shown in Fig. 2. We remark that this L changes very little in the region $15^\circ < \gamma_0 < 30^\circ$.

Intraband B(E2) transition rates of the wobbling bands, as well as interband B(E2) transition rates among wobbling bands are shown in Table 2, again for $\gamma_0 = 15^\circ$ and $\gamma_0 = 20^\circ$, normalized to the B(E2) transition rate between the two lowest $n_w = 0$ states. Some clear features are observed:

1) Intraband $L + 2 \rightarrow L$ transitions are strong, while intraband $L \rightarrow L$ transitions are weak. For interband $(n_w = 2) \rightarrow (n_w = 0)$ transitions the opposite situation appears, i.e. they are strong for $L \rightarrow L$ and weak for $L + 2 \rightarrow L$.

2) Interband $(n_w = 1) \rightarrow (n_w = 0)$ transitions are strong for $L \rightarrow L + 1$ and weak for $L \rightarrow L - 1$, while for interband $(n_w = 1) \rightarrow (n_w = 2)$ transitions the opposite picture appears, i.e. they are strong for $L \rightarrow L - 1$ and weak for $L \rightarrow L + 1$.

It is worth comparing these results to the main features expected to be exhibited by B(E2)s in wobbling bands [21].

1) In region 2 of the Lund convention, which corresponds to the present case, the interband $(n_w = 1) \rightarrow (n_w = 0)$ transitions are expected to be strong for $L \rightarrow L + 1$ and weak for $L \rightarrow L - 1$ [7]. This is exactly the situation seen in Table 2.

2) The ratio

$$\frac{B(E2)_{out}}{B(E2)_{in}} = \frac{B[E2; L_1 \rightarrow (L + 1)_0]}{B[E2; L_1 \rightarrow (L - 2)_1]}, \quad (34)$$

where the notation L_{n_w} is used, is expected [21] to be of the order 0.2–0.3, i.e. much larger than what is expected for typical interband transitions. The $(n_w = 1) \rightarrow (n_w = 0)$ transitions in Table 2 do exhibit this behaviour.

3) The $B(E2)_{out} = B[E2; L_1 \rightarrow (L + 1)_0]$ values are expected to go as $1/L$ and not as $1/L^2$ [21]. The results in Table 2 do exhibit this feature.

6. Comparison to experiment

The results of the present approach should be compared to experimental data for nuclei which exhibit a X(5) ground state band. Rare earth nuclei with this property have been summarized in Ref. [22]. In Table 3 we see that the gsb of ^{156}Dy [23] is in good agreement with the X(5) predictions, while two additional bands (the K and M bands of Ref. [23]) respectively) correspond very well to the $n_w = 0$ and $n_w = 1$ W(5) bands with $\gamma_0 = 20^\circ$, although no rms fitting with respect to γ_0 has been performed. It is therefore of great interest to measure intraband and interband B(E2) transitions for these bands, in order to see if they will exhibit the characteristic wobbling features mentioned in Section 5. It should be noticed that the spin assignments in the M band are based on the only transition seen experimentally [23, 28] to connect a level of this band to the $L = 30$ level of the K band. Since in the $(n_w = 1) \rightarrow (n_w = 0)$ case the transitions $L \rightarrow L + 1$ are the strong ones,

as found in Section 5, it is assumed that the level of the M band from which this transition starts is the $L = 29$ level.

It should be remembered at this point that in general γ -soft models involving γ -fluctuations and γ -rigid models corresponding to large rigid triaxiality lead to similar results for many observables [21, 29], when γ_{rms} of the former equals γ_{rigid} of the latter. Therefore the agreement of the present model to experiment, if proved for B(E2)s, too, offers indeed evidence for triaxiality but not necessarily for γ -rigid behaviour, although γ -rigidity was among the assumptions which led to this model.

7. Discussion

In summary, a W(5) model describing the wobbling bands coexisting with a X(5) ground state band in even nuclei has been introduced. Separation of variables is achieved by assuming that the potential has a sharp minimum at $\gamma = \gamma_0$. The model predictions for given value of γ_0 are parameter-free (up to overall scale factors). The W(5) predictions for wobbling bands for $\gamma_0 = 20^\circ$ are in good agreement with experimental spectra for ^{156}Dy , the ground state band of which is described satisfactorily by X(5), while the W(5) predictions for intraband and interband B(E2) transition probabilities exhibit the features expected for wobbling bands. A characteristic feature of the model is that the $n_w = 0$ wobbling band is not coinciding with the gsb, but with the superb band crossing the gsb.

Concerning further work, the following comments can be made:

1) There exist nuclei (^{160}Yb , ^{158}Er), the ground state bands of which are described well by the X(5)- β^4 and X(5)- β^6 models [30] respectively. These models correspond to the use of β^4 and β^6 potentials in the X(5) framework, leading to $R_4 = E(4)/E(2)$ ratios of 2.769 and 2.824 respectively. It is worth using the β^4 and β^6 potentials in the W(5) framework as well, in order to examine if the parameter-free (up to γ_0) predictions for wobbling bands which will occur in these models agree with experiment.

2) The β -equation [Eq. (11)] obtained above in the W(5) framework is also exactly soluble [31, 32] for the Davidson potentials [33]

$$u(\beta) = \beta^2 + \frac{\beta_0^4}{\beta^2}, \quad (35)$$

where β_0 is the position of the minimum of the potential. In analogy to earlier work in the E(5) and X(5) frameworks [34] it is expected that $\beta_0 = 0$ will correspond to a “wobbling vibrator”, while $\beta_0 \rightarrow \infty$ will lead to the original wobbling rotator of Ref. [1].

3) Using the variational procedure developed recently in the E(5) and X(5) frameworks [34], one should be able to prove that the W(5) model can be obtained from the Davidson potentials by maximizing the rate of change of various measures of collectivity with respect to the parameter β_0 , thus proving that W(5) corresponds to the critical point symmetry of the transition from a “wobbling vibrator” to a wobbling rotator.

Work in these directions is in progress.

Acknowledgements

The authors are thankful to Jean Libert (Orsay) and Werner Scheid (Giessen) for illuminating discussions.

References

- [1] A. Bohr and B. R. Mottelson, Nuclear Structure, Benjamin, New York, 1975, Vol. II, section 4-5e.
- [2] S. W. Ødegård et al., Phys. Rev. Lett. 86 (2001) 5866.
- [3] D. R. Jensen et al., Phys. Rev. Lett. 89 (2002) 142503.
- [4] D. R. Jensen et al., Nucl. Phys. A 703 (2002) 3.
- [5] G. Schönwaßer et al., Phys. Lett. B 552 (2003) 9.
- [6] H. Amro et al., Phys. Lett. B 553 (2003) 197.
- [7] Y. R. Shimizu and M. Matsuzaki, Nucl. Phys. A 588 (1995) 559.
- [8] M. Matsuzaki, Y. R. Shimizu, and K. Matsuyanagi, Phys. Rev. C 65 (2002) 041303.
- [9] M. Matsuzaki, Y. R. Shimizu, and K. Matsuyanagi, Phys. Rev. C 69 (2004) 034325.
- [10] I. Hamamoto, Phys. Rev. C 65 (2002) 044305.
- [11] I. Hamamoto, Phys. Rev. C 67 (2003) 014319.
- [12] F. Iachello, Phys. Rev. Lett. 85 (2000) 3580.
- [13] F. Iachello, Phys. Rev. Lett. 87 (2001) 052502.
- [14] F. Iachello, Phys. Rev. Lett. 91 (2003) 132502.
- [15] D. Bonatsos, D. Lenis, D. Petrellis, and P. A. Terziev, Phys. Lett. B, in press. nucl-th/0402087.
- [16] A. S. Davydov and G. F. Filippov, Nucl. Phys. 8 (1958) 237.
- [17] A. S. Davydov and V. S. Rostovsky, Nucl. Phys. 12 (1959) 58.
- [18] A. S. Davydov, Nucl. Phys. 24 (1961) 682.
- [19] A. Bohr, Mat. Fys. Medd. K. Dan. Vidensk. Selsk. 26 (1952) no. 14.
- [20] R. J. Turner and T. Kishimoto, Nucl. Phys. A 217 (1971) 317.
- [21] R. F. Casten, E. A. McCutchan, N. V. Zamfir, C. W. Beausang, and J.-Y. Zhang, Phys. Rev. C 67 (2003) 064306.
- [22] E. A. McCutchan et al., Phys. Rev. C 69 (2004) 024308.
- [23] C. W. Reich, Nucl. Data Sheets 99 (2003) 753.
- [24] R. Bijker, R. F. Casten, N. V. Zamfir, and E. A. McCutchan, Phys. Rev. C 68 (2003) 064304.

- [25] J. Meyer-ter-Vehn, Nucl. Phys. A 249 (1975) 111.
- [26] A. R. Edmonds, Angular Momentum in Quantum Mechanics, Princeton University Press, Princeton, 1957.
- [27] T. Bengtsson, Nucl. Phys. A 496 (1989) 56; 512 (1990) 124.
- [28] F. G. Kondev et al., Phys. Lett. B 437 (1998) 35.
- [29] R. F. Casten, Nuclear Structure from a Simple Perspective, Oxford University Press, Oxford, 1990.
- [30] D. Bonatsos, D. Lenis, N. Minkov, P. P. Raychev, and P. A. Terziev, Phys. Rev. C 69 (2004) 014302.
- [31] J. P. Elliott, J. A. Evans, and P. Park, Phys. Lett. B 169 (1986) 309.
- [32] D. J. Rowe and C. Bahri, J. Phys. A 31 (1998) 4947.
- [33] P. M. Davidson, Proc. R. Soc. 135 (1932) 459.
- [34] D. Bonatsos, D. Lenis, N. Minkov, D. Petrellis, P. P. Raychev, and P. A. Terziev, Phys. Lett. B 584 (2004) 40.

Figure captions

Fig. 1 Energy levels with various L (and $s = 1$) of the W(5) model plotted versus γ_0 , the position at which the potential has a sharp minimum. a) The $n_w = 0$ band. b) The $n_w = 1$ band. See Section 5 for further discussion.

Fig. 2 Angular momentum L up to which the X(5) ground state band is Yrast, while beyond it the $n_w = 0$ W(5) band becomes Yrast, as a function of γ_0 , the position at which the potential has a sharp minimum. See Section 5 for further discussion.

Table 1: Energy levels of the W(5) model (with $s = 1$), measured from the ground state of X(5) and normalized to the first excited state of X(5). The ground state band of X(5), normalized in the same way, is shown for comparison. See Sections 2 and 5 for further details.

	X(5)	W(5)	W(5)	W(5)	W(5)		W(5)	W(5)	W(5)	W(5)
n_w		0	0	2	2		1	1	3	3
γ_0		15°	20°	15°	20°		15°	20°	15°	20°
L_0		2.061	1.378	10.307	6.891		6.184	4.134	14.429	9.647
L						L				
0	0.000	(0.000)	(0.000)							
2	1.000	(1.614)	1.513	(4.293)	(3.967)	3	(4.535)	(4.222)		
4	2.904	3.729	3.531	(8.276)	(7.692)	5	(7.744)	7.253	(12.766)	(11.850)
6	5.430	6.265	5.964	(12.427)	(11.593)	7	11.257	10.586	(17.790)	(16.555)
8	8.483	9.195	8.780	(16.841)	15.755	9	15.099	14.241	(23.062)	(21.506)
10	12.027	12.504	11.964	(21.555)	20.210	11	19.281	18.226	(28.618)	26.734
12	16.041	16.184	15.507	26.585	24.973	13	23.806	22.543	(34.475)	32.256
14	20.514	20.227	19.401	31.940	30.052	15	28.675	27.194	40.645	38.082
16	25.437	24.630	23.642	37.624	35.450	17	33.887	32.177	47.134	44.218
18	30.804	29.389	28.226	43.641	41.170	19	39.444	37.492	53.945	50.667
20	36.611	34.501	33.151	49.991	47.212	21	45.343	43.138	61.083	57.431
22	42.853	39.962	38.413	56.676	53.577	23	51.584	49.113	68.548	64.511
24	49.528	45.772	44.011	63.695	60.266	25	58.166	55.419	76.342	71.910
26	56.633	51.927	49.942	71.049	67.278	27	65.087	62.052	84.464	79.626
28	64.166	58.427	56.205	78.737	74.613	29	72.348	69.011	92.917	87.661
30	72.124	65.270	62.798	86.759	82.270	31	79.947	76.297	101.699	96.014
32	80.505	72.454	69.720	95.116	90.250	33	87.883	83.908	110.811	104.686
34	89.309	79.978	76.970	103.805	98.551	35	96.156	91.843	120.253	113.675
36	98.533	87.841	84.546	112.828	107.173	37	104.765	100.102	130.024	122.983
38	108.176	96.042	92.448	122.183	116.115	39	113.708	108.684	140.126	132.609
40	118.237	104.580	100.675	131.870	125.378	41	122.986	117.588	150.556	142.552
42	128.715	113.454	109.225	141.888	134.960	43	132.597	126.813	161.315	152.813
44	139.608	122.664	118.098	152.238	144.862	45	142.541	136.360	172.404	163.391
46	150.917	132.208	127.294	162.919	155.082	47	152.818	146.226	183.821	174.285
48	162.639	142.086	136.810	173.930	165.621	49	163.427	156.412	195.566	185.496
50	174.774	152.297	146.648	185.270	176.477	51	174.367	166.917	207.639	197.023
52	187.321	162.840	156.806	196.941	187.651	53	185.637	177.741	220.040	208.867
54	200.280	173.715	167.284	208.940	199.142	55	197.238	188.884	232.768	221.025
56	213.650	184.922	178.080	221.268	210.950	57	209.169	200.344	245.823	233.499
58	227.430	196.459	189.195	233.925	223.074	59	221.430	212.121	259.206	246.288
60	241.620	208.327	200.629	246.909	235.514	61	234.019	224.215	272.915	259.391

Table 2: B(E2) transition rates of the W(5) model (connecting states with $s = 1$), normalized to the transition between the two lowest states, B(E2; $2_0 \rightarrow 0_0$). See Sections 4 and 5 for further details.

$L_{n_w}^{(i)}$	γ_0 $L_{n_w}^{(f)}$	15°	20°	$L_{n_w}^{(i)}$	$L_{n_w}^{(f)}$	15°	20°	$L_{n_w}^{(i)}$	$L_{n_w}^{(f)}$	15°	20°
2 ₀	0 ₀	1.000	1.000								
4 ₀	2 ₀	1.606	1.597	4 ₂	2 ₂	0.785	0.777	5 ₁	3 ₁	1.268	1.256
6 ₀	4 ₀	2.228	2.213	6 ₂	4 ₂	1.072	1.061	7 ₁	5 ₁	1.889	1.872
8 ₀	6 ₀	2.665	2.647	8 ₂	6 ₂	1.635	1.620	9 ₁	7 ₁	2.349	2.330
10 ₀	8 ₀	2.998	2.980	10 ₂	8 ₂	2.081	2.063	11 ₁	9 ₁	2.705	2.685
12 ₀	10 ₀	3.265	3.247	12 ₂	10 ₂	2.440	2.421	13 ₁	11 ₁	2.992	2.971
14 ₀	12 ₀	3.485	3.468	14 ₂	12 ₂	2.734	2.715	15 ₁	13 ₁	3.228	3.208
16 ₀	14 ₀	3.672	3.655	16 ₂	14 ₂	2.981	2.962	17 ₁	15 ₁	3.429	3.409
18 ₀	16 ₀	3.832	3.816	18 ₂	16 ₂	3.192	3.173	19 ₁	17 ₁	3.601	3.582
20 ₀	18 ₀	3.972	3.956	20 ₂	18 ₂	3.374	3.356	21 ₁	19 ₁	3.752	3.734
2 ₀	2 ₀	0.114	0.049	2 ₂	2 ₂	0.131	0.056				
4 ₀	4 ₀	0.228	0.098	4 ₂	4 ₂	0.021	0.009	5 ₁	5 ₁	0.047	0.020
6 ₀	6 ₀	0.307	0.132	6 ₂	6 ₂	0.003	0.001	7 ₁	7 ₁	0.118	0.051
8 ₀	8 ₀	0.366	0.158	8 ₂	8 ₂	0.036	0.016	9 ₁	9 ₁	0.184	0.079
10 ₀	10 ₀	0.413	0.178	10 ₂	10 ₂	0.084	0.036	11 ₁	11 ₁	0.240	0.103
12 ₀	12 ₀	0.451	0.194	12 ₂	12 ₂	0.132	0.057	13 ₁	13 ₁	0.288	0.124
14 ₀	14 ₀	0.482	0.208	14 ₂	14 ₂	0.178	0.076	15 ₁	15 ₁	0.329	0.142
16 ₀	16 ₀	0.510	0.220	16 ₂	16 ₂	0.219	0.094	17 ₁	17 ₁	0.365	0.157
18 ₀	18 ₀	0.533	0.230	18 ₂	18 ₂	0.257	0.111	19 ₁	19 ₁	0.397	0.171
20 ₀	20 ₀	0.554	0.239	20 ₂	20 ₂	0.291	0.125	21 ₁	21 ₁	0.424	0.183
2 ₂	2 ₀	1.657	1.647	3 ₁	4 ₀	1.267	1.256	3 ₁	2 ₂	2.292	2.267
4 ₂	4 ₀	0.352	0.350	5 ₁	6 ₀	0.988	0.980	5 ₁	4 ₂	1.355	1.341
6 ₂	6 ₀	0.199	0.198	7 ₁	8 ₀	0.819	0.814	7 ₁	6 ₂	1.290	1.279
8 ₂	8 ₀	0.130	0.129	9 ₁	10 ₀	0.705	0.700	9 ₁	8 ₂	1.187	1.177
10 ₂	10 ₀	0.092	0.092	11 ₁	12 ₀	0.621	0.617	11 ₁	10 ₂	1.086	1.078
12 ₂	12 ₀	0.069	0.069	13 ₁	14 ₀	0.556	0.554	13 ₁	12 ₂	0.998	0.991
14 ₂	14 ₀	0.054	0.054	15 ₁	16 ₀	0.505	0.503	15 ₁	14 ₂	0.921	0.916
16 ₂	16 ₀	0.043	0.043	17 ₁	18 ₀	0.463	0.461	17 ₁	16 ₂	0.855	0.851
18 ₂	18 ₀	0.036	0.036	19 ₁	20 ₀	0.428	0.426	19 ₁	18 ₂	0.798	0.794
20 ₂	20 ₀	0.030	0.030	21 ₁	22 ₀	0.398	0.397	21 ₁	20 ₂	0.748	0.744
2 ₂	0 ₀	0.072	0.031	3 ₁	2 ₀	0.149	0.064	3 ₁	4 ₂	0.164	0.070
4 ₂	2 ₀	0.049	0.021	5 ₁	4 ₀	0.152	0.065	5 ₁	6 ₂	0.216	0.093
6 ₂	4 ₀	0.029	0.012	7 ₁	6 ₀	0.140	0.060	7 ₁	8 ₂	0.220	0.095
8 ₂	6 ₀	0.019	0.008	9 ₁	8 ₀	0.127	0.055	9 ₁	10 ₂	0.212	0.091
10 ₂	8 ₀	0.013	0.006	11 ₁	10 ₀	0.113	0.050	11 ₁	12 ₂	0.200	0.086
12 ₂	10 ₀	0.010	0.004	13 ₁	12 ₀	0.107	0.046	13 ₁	14 ₂	0.188	0.081
14 ₂	12 ₀	0.008	0.003	15 ₁	14 ₀	0.098	0.042	15 ₁	16 ₂	0.176	0.076
16 ₂	14 ₀	0.006	0.003	17 ₁	16 ₀	0.091	0.039	17 ₁	18 ₂	0.166	0.071

Table 3: Comparison of the X(5) predictions for the ground state band and the W(5) predictions (with $\gamma_0 = 20^\circ$) for the $n_w = 0, 1$ bands (with $s = 1$) to experimental data for ^{156}Dy [23]. See Section 6 for further discussion.

n_w	X(5)		exp.	W(5)		exp.	W(5)		exp.	W(5)		exp.
	0			0			1					
L	L	L	L	L	L	L	L	L	L	L	L	
0	0.000	0.000	14	19.401	22.254	38	92.448	92.686	29	69.011	70.349	
2	1.000	1.000	16	23.642	25.396	40	100.675	100.785	31	76.297	77.150	
4	2.904	2.934	18	28.226	29.221	42	109.225	108.839	33	83.908	84.300	
6	5.430	5.592	20	33.151	33.647	44	118.098	117.378	35	91.843	91.660	
8	8.483	8.823	22	38.413	38.617	46	127.294	125.920	37	100.102	99.339	
10	12.027	12.521	24	44.011	44.060	48	136.810	135.116	39	108.684	107.425	
12	16.041	16.592	26	49.942	49.923	50	146.648	144.828	41	117.588	115.954	
14	20.514	20.961	28	56.205	56.172	52	156.806	155.491	43	126.813	125.107	
16	25.437	25.574	30	62.798	62.792	54	167.284	166.930	45	136.360	135.008	
18	30.804	30.327	32	69.720	69.763	56	178.080	179.400	47	146.226	145.184	
20	36.611	35.269	34	76.970	77.070	58	189.195	193.366	49	156.412	156.144	
22	42.853	40.451	36	84.546	84.711				51	166.917	168.716	





